



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Erythea, these last at the base of the mountains. In the southern third of the peninsula many distinctly tropical genera appear, such as *Ficus*, *Mimosa*, *Cassia*, *Albizzia*, *Jatropha*, *Haematoxylon*, *Lantana*, *Manihot*, and *Chiococca*.

Among the more remarkable endemic desert forms, two trees may be mentioned. One, belonging to the Anacardiaceae, *Pachycormus discolor*, is found in the extremely arid central section of the peninsula. Seldom 10 ft. in height, the branches often shoot out twice that distance from the trunk, while their thickness (1 ft. or more), their abrupt ending in a few short twigs covered with red flowers, "reminding one of the proboscis of an elephant holding a nosegay," give a remarkably grotesque appearance to the tree. The leaves are minute and fall off before the flowers are fully developed. The associated monotypic genus *Idria columnaris* is in the Fouquieriaceae, and appears as a tree reaching 50 ft. in height in a scattered open forest. In contrast with the preceding it has a straight columnar trunk, usually without large branches. Illustrations of these and many other interesting and unusual plants add much to the interest of both reports.—GEO. D. FULLER.

Rubus in New England.—BRAINERD and PEETERSEN,²¹ recognizing that "*Rubus* is one of the most polymorphic genera in the entire plant kingdom," have presented the blackberry group of that genus as displayed in New England. The authors say that the remarkable variation in the number of species recognized in the various taxonomic works is due to too great reliance upon herbarium specimens, to failure to appreciate the variations due to environmental conditions, and to lack of appreciation of the extent of interbreeding. The present study is based upon data from material in the field, behavior in garden cultures and controlled plots, characters of the progeny of supposed natural hybrids, and behavior of progeny when artificially crossed. The result is that the authors recognize twelve valid species of New England blackberries, and a long list of hybrids.

In following up the experimental work, PEETERSEN²² has reached the following conclusions: variations due to external factors are very marked; primordia of the prickle, glandular hair, and simple hair are present in all species; a large percentage of infertility occurs in most species, largely due to defective pollen; cross pollination is the rule in all species, all the species studied being either nearly or completely self-sterile; all the species are capable of inter-crossing under favorable conditions; duplicates of natural hybrids were produced artificially; the progeny of a number of so-called species segregated as hybrids.

The paper is a good illustration of the test of genetics applied to taxonomy.—J. M. C.

²¹ BRAINERD, EZRA, and PEETERSEN, A. K., Blackberries of New England; their classification. Bull. 217, Vermont Agric. Exper. Sta. pp. 84. pls. 36. 1920.

²² PEETERSEN, A. K., Blackberries of New England; genetic status of the plants. Bull. 218, Vermont Agric. Exper. Sta. pp. 34. pls. 19. 1921.